

## Differentiation

Recall that  $f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$  is the *derivative of  $f$  at  $a$*  (provided the limit exists). To avoid problems we take  $x$  to be in an *open* interval on which  $f$  is defined. The derivative function  $f'$  is the function whose value at  $a$  is  $f'(a)$ .

There are classical examples of continuous but nowhere differentiable functions. However in the other direction we have the following result:

**Theorem**  $f$  differentiable at  $a \implies f$  continuous at  $a$ .

Rules for derivatives of sums and products are just particular cases of previous results about limits. Higher derivatives are defined recursively, i.e.  $f^{(n)} = (f^{(n-1)})'$ . Most common functions are infinitely differentiable (of class  $C^\infty$ ) except possibly at a finite or countable set of points.

**Chain Rule** If  $f$  is differentiable at  $x$  and  $g$  is differentiable at  $f(x)$  then the composite function  $g \circ f$  is differentiable at  $x$  and

$$(g \circ f)'(x) = g'[f(x)]f'(x).$$

**Proof** Note first that  $f$  is differentiable at  $x$  if and only if there is a number  $L$  and a function  $E(x, h)$  such that

$$f(x+h) = f(x) + Lh + hE(x, h), \quad \text{where } E(x, h) \rightarrow 0 \text{ as } h \rightarrow 0.$$

In fact if  $f$  is differentiable at  $x$  we can take  $E(x, h) = \frac{f(x+h) - f(x)}{h} - f'(x)$  and we can make  $E(x, h)$  continuous at 0 by defining  $E(x, 0) = 0$ .

Now let  $k = k(h) = f(x+h) - f(x)$  and let  $y = f(x)$ . Then

$$g[f(x+h)] - g[f(x)] = g(y+k) - g(y) = g'(y)k + kE(y, k),$$

where  $E(y, k) \rightarrow 0$  as  $k \rightarrow 0$  (and  $E(y, 0) = 0$  by definition). So

$$\frac{g[f(x+h)] - g[f(x)]}{h} = g'[f(x)] \frac{f(x+h) - f(x)}{h} + \frac{f(x+h) - f(x)}{h} E[f(x), k(h)].$$

Now take limits as  $h \rightarrow 0$  and use the fact that the functions  $E$  and  $k$  are continuous at 0 and take the value 0 there.

The following result is crucial in the application of differential calculus to maximum and minimum problems:

**Theorem** Let  $f$  be defined on an open interval  $I$ , and assume that  $f$  has a relative maximum or minimum at an *interior* point  $p$  of  $I$ . If the derivative  $f'(p)$  exists then  $f'(p) = 0$ .

**Proof** Define a function  $Q$  on  $I$  by  $Q(x) = \frac{f(x) - f(p)}{x - p}$  for  $x \neq p$  and  $Q(p) = f'(p)$ . Since  $f'(p)$  exists,  $Q(x) \rightarrow Q(p)$  as  $x \rightarrow p$  and so  $Q$  is continuous at  $p$ . We must show that  $Q(p) = 0$ .

Suppose  $Q(p) > 0$ . By the sign-preserving property of continuous functions there is an interval about  $p$  in which  $Q(x)$  is positive. So the numerator and denominator of  $Q(x)$  have the same sign for all  $x \neq 0$  in this interval, i.e.

$$f(x) > f(p) \text{ for } x > p \quad \text{and} \quad f(x) < f(p) \text{ for } x < p.$$

But this contradicts the assumption that  $f$  has an extremum at  $p$ . Hence  $Q(p) > 0$  is impossible. Similarly  $Q(p) < 0$  is impossible. So  $Q(p) = 0$ .

**Rolles Theorem** If  $f$  is continuous on  $[a, b]$ , differentiable on  $(a, b)$  and  $f(a) = f(b)$ , then there is at least one point  $p$  in  $(a, b)$  such that  $f'(p) = 0$ .

**Proof** Since  $f$  is continuous on  $[a, b]$  it follows from the Extreme Value Theorem that it has a maximum and a minimum in  $[a, b]$ . If there is a maximum  $M$  or minimum  $m$  at  $p \in (a, b)$  then  $f'(p) = 0$  by the previous theorem. If both extreme values occur at endpoints then  $M = m$  since  $f(a) = f(b)$  and so  $f$  is constant on  $[a, b]$ , i.e.  $f'(p) = 0$  for any  $p \in (a, b)$ .

**Mean Value Theorem** If  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  then there is at least one point  $p \in (a, b)$  such that  $f(b) - f(a) = (b - a)f'(p)$ .

**Proof** We apply Rolles Theorem to the function

$$g(x) = \left[ \frac{f(b) - f(a)}{b - a} \right] (x - a) + f(a) - f(x).$$

### Remark

From the Mean Value Theorem we can deduce the standard results about the geometric interpretation of derivatives (e.g.  $f$  is strictly increasing on  $[a, b]$  if  $f'(x) > 0$  for all  $x \in (a, b)$ ) and thus the first and second derivative tests for extrema.

Another consequence is that  $f$  is constant on  $[a, b]$  if  $f'(x) = 0$  on  $(a, b)$ . It follows that if  $g$  is another function with the same properties as  $f$  and  $f'(x) = g'(x)$  for all  $x \in (a, b)$  then  $f$  and  $g$  differ only by a constant. This is a crucial point in showing that we can calculate integrals in terms of antiderivatives.